

GENERATION OF SURFACE COORDINATES BY ELLIPTIC

PARTIAL DIFFERENTIAL EQUATIONS*

Z.U.A. Warsi

Department of Aerospace Engineering
Mississippi State University
Mississippi State, MS 39762

The problem of generating spatial coordinates by numerical methods through carefully selected mathematical models is of current interest both in mechanics and physics. A review of various methods of coordinate generation in two and three-dimensional Euclidean spaces is available in reference 1, and reference may also be made to the proceedings of two recent conferences (references 2 and 3) and a book (ref. 4) on the topic of numerical grid generation.

In this paper the problem of generation of a desired system of coordinates in a given surface has been considered which essentially is an effort directed to the problem of grid generation in a two-dimensional non-Euclidean space. The mathematical model selected for this purpose is based on the formulae of Gauss for a surface and has been discussed by the author in earlier publications (refs. 5 - 9).

The formulae of Gauss for a surface $v=\text{const.}$ can be written compactly by using the summation convention on repeated lower and upper indices as

$$\underline{r}_{,\alpha\beta} = T_{\alpha\beta}^{\delta} \underline{r}_{,\delta} + \underline{n}^{(\nu)} b_{\alpha\beta} \quad (1)$$

where $\underline{r}=(x,y,z)$, $T_{\alpha\beta}^{\delta}$ are the surface Christoffel symbols, $\underline{n}^{(\nu)}$ is the unit surface normal on the surface $v=\text{const.}$, $b_{\alpha\beta}$ are the coefficients of the second fundamental form, and a comma denotes the partial derivative with respect to the surface coordinates $x^{\alpha(v)}$ and the other Greek indices assuming cyclic values). Inner multiplication by $G_{\nu} g^{\alpha\beta}$ of equation (1) yields

$$D\underline{r} + G_{\nu} (\Delta_2^{(\nu)} x^{\delta}) \underline{r}_{,\delta} = \underline{n}^{(\nu)} R, \quad (2)$$

where

$$G_{\nu} = g_{\alpha\alpha} g_{\beta\beta} - (g_{\alpha\beta})^2, \quad \nu, \alpha, \beta \text{ cyclic},$$

$$D = G_{\nu} g^{\alpha\beta} \partial_{\alpha\beta}, \quad (3a-c)$$

$$R = G_{\nu} g^{\alpha\beta} b_{\alpha\beta} = G_{\nu} (k_I^{(\nu)} + k_{II}^{(\nu)}),$$

*Research performed under grant AFOSR-0185.

and

$$\Delta_2^{(\nu)} x^\alpha = -g^{\beta\gamma} T_{\beta\gamma}^\alpha, \quad (3d)$$

is the second-order differential parameter of Beltrami. In equation (3c), $k_I^{(\nu)}$ and $k_{II}^{(\nu)}$ are the principal curvatures at a point on the surface $\nu=\text{const.}$

Equation (2) provides three coupled quasilinear elliptic partial differential equations with the Cartesian coordinates x, y, z as dependent variables. These equations are nonhomogeneous with the right hand sides depending on the components of both the normal and the mean curvature of the surface, thus reflecting some geometrical aspects of the surface in an explicit manner.

Some of the problems listed below have been solved successfully by taking equation (2) as a basic differential model.

I. If the mathematical equation of the surface is available in the form $F(x, y, z)=0$, then equation (2) can be used to introduce any desired coordinate system in the surface. (For discrete x, y, z values of a surface, the form $F(x, y, z)=0$ can be obtained either by a least square or piecewise approximation method. Knowing $F(x, y, z)=0$, one can find $k_I^{(\nu)} + k_{II}^{(\nu)}$ as a function of x, y, z . Now two options are open: In the first, one can retain $\Delta_2^{(\nu)} x^\delta$ as it appears in the equation, and in the second write $\Delta_2^{(\nu)} x^\delta = P^\delta / G_\nu$, where P^δ are arbitrarily specified functions. The second option provides a control by the user on the distribution of coordinates in the surface.

II. The proposed equations can be used to generate a new coordinate system from the data of an already given coordinate system in a surface (refs. 10 and 11).

III. If the coefficients of the first and second fundamental forms have been given, then the proposed equations can be used to generate a surface satisfying the given data (surface fitting).

IV. The proposed equations can also be used to generate surfaces in the space between two arbitrary given surfaces, thus providing 3D grids in an Euclidean space (refs. 6, 7, 12, and 13).

A number of numerical and analytical results obtained by the author and his co-workers will be presented in the seminar.

REFERENCES

1. Thompson, J.F.; Warsi, Z.U.A.; Mastin, C.W. Journal of Computational Physics 47, 1982, pp. 1-108.
2. Thompson, J.F. (Ed.), Numerical Grid Generation, North-Holland, 1982.
3. Ghia, K.N. and Ghia, U. (Ed.), Advances in Grid Generation, ASME Publication No. Fed-Vol. 5, 1983.
4. Thompson, J.F.; Warsi, Z.U.A.; and Mastin, C.W. Numerical Grid Generation: Foundations and Applications, North-Holland, 1985.
5. Warsi, Z.U.A., "A Method for the Generation of Three-Dimensional Coordinates Between Bodies of Arbitrary Shapes", MSSU-EIRS-90-7, Mississippi State University, 1980.
6. Warsi, Z.U.A., "Basic Differential Models for Coordinate Generation", in Numerical Grid Generation, Ed. J.F. Thompson, North-Holland, 1982, pp. 41-77.
7. Warsi, Z.U.A., Quart. Appl. Math., 41, 1983, pp. 221-236.
8. Warsi, Z.U.A. and Ziebarth, J.P., "Numerical Generation of Three-Dimensional Coordinates Between Bodies of Arbitrary Shapes", in Numerical Grid Generation, Ed. J.F. Thompson, North-Holland 1982, pp. 717-728.
9. Warsi, Z.U.A., "Numerical Grid Generation in Arbitrary Surfaces Through a Second-Order Differential Geometric Models", Journal of Computational Physics, submitted.
10. Tiarn, W-N., "Generation of Surface Coordinates in Curved Surfaces and in Plane Regions", M.S. Thesis, Mississippi State University, December 1983.
11. Ziebarth, J.P. and Warsi, Z.U.A., "Computer Simulation of Three-Dimensional Grids", Society of Computer Simulation, February 1984.
12. Ziebarth, J.P., "Three-Dimensional Grid Generation Through Elliptic Partial Differential Equations", Ph.D. Dissertation, Mississippi State University, December 1983.
13. Warsi, Z.U.A., "Generation of Three-Dimensional Grids Through Elliptic Differential Equation", CFD Lecture Series, Von Karman Institute for Fluid Dynamics, March 1984.